

**PHEXXI (lactic acid, citric acid, and potassium bitartrate) Vaginal Gel**  
**Important Safety Information**

**INDICATIONS AND USAGE**

PHEXXI is indicated for the prevention of pregnancy in females of reproductive potential for use as an on-demand method of contraception.

**LIMITATIONS OF USE**

PHEXXI is not effective for the prevention of pregnancy when administered after intercourse.

**WARNINGS AND PRECAUTIONS**

Few cases (0.36%) reported adverse reactions of cystitis, pyelonephritis and other upper urinary tract infection (UTI) have been reported in PHEXXI clinical studies. Of these, one case of pyelonephritis was considered serious and required hospitalization. Avoid use of PHEXXI in females of reproductive potential with history of recurrent urinary tract infection or urinary tract abnormalities .

**ADVERSE REACTIONS**

Most common adverse reactions ( $\geq 2\%$ ) were vulvovaginal burning sensation, vulvovaginal pruritus, vulvovaginal mycotic infection, urinary tract infection, vulvovaginal discomfort, bacterial vaginosis, vaginal discharge, genital discomfort, dysuria, and vulvovaginal pain.

**Patients should be counseled on the following:**

- **To contact and consult with their healthcare provider for severe or prolonged genital irritation or experiencing urinary tract symptoms.**
- **To discontinue PHEXXI if they develop a local hypersensitivity reaction.**
- **That PHEXXI does not protect against HIV infection (AIDS) or other sexually transmitted infections.**

Please see full Prescribing Information for PHEXXI.

***To report SUSPECTED ADVERSE REACTIONS, contact Evofem at toll-free phone 1-833-EVFM BIO or you may, contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).***