

What is Phexxi?

Phexxi is a prescription vaginal gel used to prevent pregnancy in females who choose to use an on-demand method of birth control. Phexxi is only effective when used immediately **before** (or up to one hour before) each act of vaginal sex. It is not effective when used after vaginal sex.

Important Safety Information**What are the possible side effects of Phexxi?**

There have been a few cases of urinary bladder infection and kidney infection reported in clinical studies. One of the cases was serious. Avoid Phexxi if you have had repeated urinary tract infections or other urinary tract problems.

The **most common side effects** were vaginal burning, vaginal itching, vaginal yeast infection, urinary tract infection, vaginal area discomfort, bacterial vaginosis, and vaginal discharge. Women also reported genital discomfort, pain while urinating, and vaginal pain. Some male partners reported genital discomfort.

What else should I know about using Phexxi?

Phexxi does not protect against any sexually transmitted diseases, including HIV. Avoid using Phexxi with a vaginal ring.

Contact your healthcare provider if you are experiencing severe genital irritation or discomfort or urinary tract symptoms. Avoid Phexxi if you or your sexual partner is allergic to lactic acid, citric acid, potassium bitartrate, or any of the ingredients in Phexxi. Stop using Phexxi if you develop an allergic reaction.

Please see full Prescribing Information for Phexxi, including Patient Information.

Please report side effects by contacting Evofem Biosciences® toll-free at 1-833-EVFM BIO or contact FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.