

Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Kimberly Daniels, Ph.D., and Joyce C. Abma, Ph.D.

Key findings


Data from the 2017–2019 National Survey of Family Growth

- In 2017–2019, 65.3% of women aged 15–49 in the United States were currently using contraception.
- The most common contraceptive methods currently used were female sterilization (18.1%), oral contraceptive pills (14.0%), long-acting reversible contraceptives (LARCs) (10.4%), and the male condom (8.4%).
- Use of LARCs was higher among women aged 20–29 (13.7%) and 30–39 (12.7%) compared with women aged 15–19 (5.8%) and 40–49 (6.6%).
- Current condom use was higher among Hispanic women (10.5%) and non-Hispanic black women (11.0%) compared with non-Hispanic white women (7.0%).
- With higher education, female sterilization declined and use of the pill increased.

Nearly all women use contraception in their lifetimes (1), although at any given time, they may not be using contraception for reasons such as seeking pregnancy, being pregnant or postpartum, or not being sexually active. Using data from the 2017–2019 National Survey of Family Growth (NSFG), this report provides a snapshot of current contraceptive status, in the month of interview, among women aged 15–49 in the United States. In addition to describing use of any method by age, Hispanic origin and race, and education, patterns of use are described for the four most commonly used contraceptive methods: female sterilization; oral contraceptive pills; long-acting reversible contraceptives (LARCs), which include contraceptive implants and intrauterine devices; and the male condom.

In 2017–2019, approximately 65% of women aged 15–49 were currently using contraception.

Figure 1. Percentage of women currently using any contraceptive method among all women aged 15–49 and by age group, Hispanic origin and race, and education: United States, 2017–2019



¹Includes persons of other and multiple race and origin groups, not shown separately.

²Significant linear trend across all four age groups.

³Significantly different from non-Hispanic black women and Hispanic women.

NOTES: The population size referenced for women aged 15–49 is 72.7 million. Analyses of education are limited to women aged 22–49 at the time of interview. Less than high school is no high school diploma or GED. Access data table for Figure 1 at: <https://www.cdc.gov/nchs/data/databriefs/db388-tables-508.pdf#1>.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics




- In 2017–2019, 65.3% of women aged 15–49 in the United States were currently using a method of contraception (Figure 1).
- Current contraceptive use increased with age, from 38.7% among women aged 15–19 to 74.8% among women aged 40–49.
- The percentage of non-Hispanic white women currently using contraception (69.2%) was higher compared with non-Hispanic black (61.4%) and Hispanic (60.5%) women.
- Current contraceptive use did not differ significantly across education (69.1%–71.3%).

Female sterilization, the pill, the male condom, and LARCs were the most common methods women reported currently using in 2017–2019.

- The most common contraceptive methods currently used among women aged 15–49 were female sterilization (18.1%), the pill (14.0%), LARCs (10.4%), and the male condom (8.4%) (Figure 2).
- In 2017–2019, approximately 35% of women aged 15–49 were not currently using contraception, including for reasons such as seeking pregnancy, being pregnant or postpartum, or not being sexually active.

Figure 2. Percent distribution of women aged 15–49, by current contraceptive status: United States, 2017–2019



¹Additional categories of nonusers, such as nonsurgical sterility, are shown in the accompanying data table.

²Other methods grouped in this category are shown in the accompanying data table.


NOTES: Percentages may not add to 100 due to rounding. Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. Access data table for Figure 2 at: <https://www.cdc.gov/nchs/data/databriefs/db388-tables-508.pdf#2>.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Current use of female sterilization, the pill, the male condom, and LARCs varied by age.

- Among women aged 20 and over, female sterilization increased with increasing age (Figure 3). Roughly 2 in 5 women aged 40–49 (39.1%) were relying on female sterilization for contraception compared with 1 in 5 women aged 30–39 (21.2%).
- Pill use generally decreased with increasing age: 19.5% of women aged 15–19 and 21.6% of women aged 20–29 were currently using the pill compared with 10.9% of those aged 30–39, and 6.5% of women aged 40–49.
- Condom use was similar among women aged 20–29 (10.4%) and 30–39 (9.7%); use was lower among women aged 15–19 (5.1%) and 40–49 (6.5%). The observed difference between women aged 15–19 and those aged 40–49 was not statistically significant.
- Use of LARCs was similar among women aged 20–29 (13.7%) and 30–39 (12.7%). Compared with these age groups, use was lower among women aged 15–19 (5.8%) and 40–49 (6.6%).

Figure 3. Percentage of all women aged 15–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by age group: United States, 2017–2019



* Percentage does not meet National Center for Health Statistics standards of reliability.

¹Significant linear trend.

²Percentages for age groups 20–29 and 30–39 were higher than percentages for age groups 15–19 and 40–49. Differences between percentages for other age groups were not statistically significant.

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. Access data table for Figure 3 at:


<https://www.cdc.gov/nchs/data/databriefs/db388-tables-508.pdf#3>.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Current use of the pill and the male condom varied by Hispanic origin and race.

- Female sterilization did not differ significantly across Hispanic-origin and race groups: 19.9% among Hispanic, 18.5% among non-Hispanic white, and 17.6% among non-Hispanic black women (Figure 4).
- Current use of the pill was higher among non-Hispanic white women (17.8%) than among Hispanic (7.9%) and non-Hispanic black (8.1%) women.
- Current condom use was lower among non-Hispanic white women (7.0%) compared with Hispanic (10.5%) and non-Hispanic black (11.0%) women.
- Current use of LARCs did not significantly differ across the Hispanic-origin and race groups shown, around 10%–11%.

Figure 4. Percentage of all women aged 15–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by Hispanic origin and race: United States, 2017–2019



¹Percentage for non-Hispanic white women is significantly different from percentages for Hispanic and non-Hispanic black women.

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. Access data table for Figure 4 at:


<https://www.cdc.gov/nchs/data/databriefs/db388-tables-508.pdf#4>.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Current use of female sterilization, the pill, and LARCs varied by educational attainment while male condom use did not.

- Among women aged 22–49, female sterilization declined with higher education, from 39.9% among women without a high school diploma or GED to 12.1% of women with a bachelor’s degree or higher (Figure 5).
- Current pill use increased with higher education: 18.1% of women with a bachelor’s degree or higher were using the pill compared with 5.7% of women without a high school diploma or GED.
- Current male condom use was similar across education, around 9%.
- Current use of LARCs was higher for women with some college but no bachelor’s degree (12.5%) and women with a bachelor’s degree or higher (13.1%) compared with women with a high school diploma or GED (7.9%), but was not significantly different from use among women without a high school diploma or GED (9.3%).

Figure 5. Percentage of all women aged 22–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by educational attainment: United States, 2017–2019



¹Significant linear trend.

²Percentages for some college, no bachelor’s degree and bachelor’s degree or higher are significantly different from that for high school diploma or GED.

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. Access data table for Figure 5 at:

<https://www.cdc.gov/nchs/data/databriefs/db388-tables-508.pdf#5>.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Summary

In 2017–2019, approximately 65% of women aged 15–49 were using some type of contraceptive method, while 35% were not currently using a method. Current use of any method of contraception was higher among older women and non-Hispanic white women compared with younger women and Hispanic and non-Hispanic black women. The most commonly used methods were female sterilization (18.1%), oral contraceptive pills (14.0%), LARCs (10.4%), and the male condom (8.4%). Use of these methods varied across age, Hispanic origin and race, and education.

Understanding variation in contraceptive use across social and demographic characteristics offers potential insight into larger fertility patterns, including birth rates and incidence of unintended pregnancies. The chance that a woman not seeking a pregnancy will have an unintended pregnancy varies by whether any method of contraception is used and which method she or her partner uses (2).

Definitions

Age: The respondent’s age at the time of the interview. The recode variable used was AGER.

Current contraceptive status: Measures contraceptive status in the month of the interview (not at a specific act of sexual intercourse). This variable includes either use of specific methods or, if the woman did not use a method in the month of interview, nonuse of contraceptive methods using the following subcategories: pregnant, seeking pregnancy, postpartum, noncontraceptive sterility, sexual inactivity (ever or in the past 3 months), or sexually active in the past 3 months and no method used in the month of interview. In this report, as in prior NSFG reports, women who were currently using more than one method are classified by the method that was most effective in preventing pregnancy, because that method has the greatest impact on their risk of unintended pregnancy (2). For example, women who report using both oral contraceptive pills and male condoms in the current month are classified as using pills, because pills are more effective at preventing pregnancy. In 2017–2019, 11.2% of women who were currently using contraception used more than one contraceptive method during the same month. The recode variable used was CONSTAT1. Although not done in this report, this set of recodes (CONSTAT1–CONSTAT4 for each method used in the month of interview) also allows for analyses of any use of each specific method.

Education: Educational attainment at the time of interview, indicating the highest degree earned or grade finished. Results by education are presented only for women aged 22 and over, as many younger women have not completed their education (79.5% of women aged 15–21 were in school at the time of interview, analysis not shown). The recode variable used was HIEDUC.

Hispanic origin and race: The 1997 Office of Management and Budget guidelines for the presentation of race and Hispanic-origin data in federal statistics are used for these classifications (3). The guidelines allowing respondents to report more than one race or ethnic origin are followed in the recode variable, HISPRACE2. Given the diversity of women categorized as non-Hispanic other single and multiple race, those results are not shown here, and the report presents results only for women who are Hispanic, non-Hispanic white single race, and non-Hispanic black single race.

Data source and methods

This report is primarily based on data from the 6,141 women in the female respondent file of the 2017–2019 NSFG and provides an update of an earlier 2015–2017 NSFG report on current contraceptive status (4). Among other information, NSFG provides detailed information about contraceptive use among U.S. women aged 15–49. The fieldwork plan, interview content, and other survey procedures are similar to previous surveys (5). NSFG is jointly planned and funded by the National Center for Health Statistics (NCHS) and several other programs of the U.S. Department of Health and Human Services. All estimates in this report are weighted to reflect the female household population aged 15–49 of the United States at the midpoint of 2017–2019 interviewing (July 2018). The response rate for the 2017–2019 NSFG was 63.4% overall, 65.2% for female respondents, and 61.4% for male respondents.

Statistics for this report were produced using the SURVEY procedures in SAS software version 9.4. Percentages were compared using two-tailed *t* tests at the 5% level. No adjustments were made for multiple comparisons. Survey clusters minus strata were used as the degrees of freedom for significance testing of pairwise comparisons. A weighted least squares regression was used to test for linear trends across age and education using the number of categories minus two as the degrees of freedom. If no linear trend was found across age and education, limited pairwise comparisons were tested. Statistical power to detect differences between groups was limited for some comparisons. The data presented in this report are bivariate associations that may be explained by other factors not controlled for in the figures or included in the report. For example, women at older ages are more likely to have completed their childbearing compared with women at younger ages. Therefore, women at younger ages would be more likely to choose nonpermanent contraceptive methods including the pill, male condom, and LARCs compared with female sterilization, a mostly permanent form of contraception (4,6). Some of the differences in contraceptive use by age, particularly use of barrier methods such as the condom, may be explained by differences in sexual activity. For example, 67.6% of females aged 15–19 did not have sex in the past 3 months compared with 20.9% of women aged 20–49 (analysis not shown). All estimates presented meet NCHS guidelines for presentation of proportions (7).

About the authors

Kimberly Daniels and Joyce C. Abma are with the National Center for Health Statistics, Division of Vital Statistics, Reproductive Statistics Branch.

**U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road, Room 4551, MS P08
Hyattsville, MD 20782–2064

FIRST CLASS MAIL
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

For more NCHS Data Briefs, visit:
<https://www.cdc.gov/nchs/products/databriefs.htm>.



NCHS Data Brief ■ No. 388 ■ October 2020

References

1. Daniels K, Mosher WD, Jones J. Contraceptive methods women have ever used: United States, 1982–2010. National Health Statistics Reports; no 62. Hyattsville, MD: National Center for Health Statistics. 2013.
2. Sundaram A, Vaughn B, Kost K, Bankole A, Finer L, Singh S, Trussell J. Contraceptive failure in the United States: Estimates from the 2006–2010 National Survey of Family Growth. *Perspect Sex Reprod Health* 49(1):7–16. 2017.
3. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Fed Regist* 62(210):58781–90. 1997. Available from: <https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>.
4. Daniels K, Abma JC. Current contraceptive status among women aged 15–49: United States, 2015–2017. NCHS Data Brief, no 327. Hyattsville, MD: National Center for Health Statistics. 2018.
5. National Center for Health Statistics. 2015–2017 National Survey of Family Growth (NSFG): Summary of design and data collection methods. Hyattsville, MD. Available from: https://www.cdc.gov/nchs/data/nsfg/PUF3-NSFG-2015-2017-Summary-of-DesignDataCollection_02Oct2019.pdf.
6. Daniels K, Daugherty J, Jones J, Mosher W. Current contraceptive use and variation by selected characteristics among women aged 15–44: United States, 2011–2013. National Health Statistics Reports; no 86. Hyattsville, MD: National Center for Health Statistics. 2015.
7. Parker JD, Talih M, Malec DJ, Beresovsky V, Carroll M, Gonzalez JF Jr, et al. National Center for Health Statistics data presentation standards for proportions. National Center for Health Statistics. *Vital Health Stat* 2(175). 2017.

Suggested citation

Daniels K, Abma JC. Current contraceptive status among women aged 15–49: United States, 2017–2019. NCHS Data Brief, no 388. Hyattsville, MD: National Center for Health Statistics. 2020.

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

National Center for Health Statistics

Brian C. Moyer, Ph.D., *Director*
Amy M. Branum, Ph.D., *Acting Associate Director for Science*

Division of Vital Statistics

Steven Schwartz, Ph.D., *Director*
Isabelle Horon, Dr.P.H., *Acting Associate Director for Science*

For e-mail updates on NCHS publication releases, subscribe online at:
<https://www.cdc.gov/nchs/email-updates.htm>.

For questions or general information about NCHS:
Tel: 1–800–CDC–INFO (1–800–232–4636)
TTY: 1–888–232–6348
Internet: <https://www.cdc.gov/nchs>
Online request form: <https://www.cdc.gov/info>

ISSN 1941–4927 Print ed.
ISSN 1941–4935 Online ed.

CS320389

Keywords: pill • condom • LARC • National Survey of Family Growth

Data Brief 388. Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Data table for Figure 1. Percentage of women currently using any contraceptive method among all women aged 15–49 and by age group, Hispanic origin and race, and education: United States, 2017–2019

Characteristic	Number (thousands)	Percent	Standard error
Total ¹	72,672	65.3	0.98
Age group			
15–19	9,439	38.7	2.27
20–29	21,392	60.9	1.74
30–39	21,581	72.3	1.51
40–49	20,260	74.8	1.45
Hispanic origin and race			
Hispanic	15,133	60.5	1.61
Non-Hispanic white	40,420	69.2	1.17
Non-Hispanic black	9,805	61.4	2.10
Education			
No high school diploma or GED	4,744	71.2	3.07
High school diploma or GED	14,530	70.0	2.34
Some college, but no bachelor's degree	17,631	71.3	1.63
Bachelor's degree or higher	22,371	69.1	1.70

¹Includes persons of other and multiple race and origin groups, not shown separately.

NOTE: Analyses of education are limited to women aged 22–49 at the time of interview.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Data Brief 388. Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Data table for Figure 2. Percent distribution of women aged 15–49, by current contraceptive status: United States, 2017–2019

Characteristic	Percent	Standard error
All women (total)	100.0	...
Not currently using contraception	34.7	0.98
Never had intercourse or no intercourse in 3 months before interview	16.8	0.71
Never had intercourse	9.9	0.50
No intercourse in 3 months before interview	6.9	0.47
Pregnant, postpartum, or seeking pregnancy	7.8	0.50
Pregnant or postpartum	3.1	0.26
Seeking pregnancy	4.8	0.36
Had intercourse in 3 months before interview	7.0	0.46
Other nonuse:	3.0	0.33
Surgically sterile—female (noncontraceptive)	0.1	0.08
Nonsurgically sterile—female or male	2.9	0.34
Currently using contraception (contraceptors)	65.3	0.98
Female sterilization	18.1	0.95
Male sterilization	5.6	0.53
Oral contraceptive pill	14.0	0.83
Male condom	8.4	0.61
Long-acting reversible contraception (LARC)	10.4	0.76
Intrauterine device (IUD)	8.4	0.64
Implant	2.0	0.27
3-month injectable (Depo-Provera), contraceptive ring, or contraceptive patch	3.1	0.31
3-month injectable (Depo-Provera)	2.0	0.23
Contraceptive ring or patch	1.1	0.20
All other methods	5.7	0.52
Diaphragm	*	...
Periodic abstinence—calendar rhythm	1.4	0.22
Periodic abstinence—safe period by temperature or cervical mucus test	0.3	0.13
Withdrawal	3.7	0.41
Other methods ¹	0.3	0.11

... Category not applicable.

* Estimate does not meet National Center for Health Statistics standards of reliability.

¹Includes emergency contraception, jelly or cream, as well as other methods.

NOTES: Percentages may not add to 100 due to rounding. Women currently using more than one method are classified according to the most effective method they are using.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Data Brief 388. Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Data table for Figure 3. Percentage of all women aged 15–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by age group: United States, 2017–2019

Age group	Number (thousands)	Female sterilization		Pill		Male condom		Long-acting reversible contraceptives	
		Percent	SE	Percent	SE	Percent	SE	Percent	SE
15–19	9,439	*	...	19.5	1.87	5.1	1.19	5.8	1.04
20–29	21,392	2.9	0.53	21.6	1.83	10.4	1.25	13.7	1.48
30–39	21,581	21.2	1.67	10.9	1.12	9.7	1.09	12.7	1.34
40–49	20,260	39.1	2.36	6.5	1.00	6.5	0.99	6.6	1.01

* Percentage does not meet National Center for Health Statistics standards of reliability.

... Category not applicable.

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. SE is standard error.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Data Brief 388. Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Data table for Figure 4. Percentage of all women aged 15–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by Hispanic origin and race: United States, 2017–2019

Hispanic origin and race	Number (thousands)	Female sterilization		Pill		Male condom		Long-acting reversible contraceptives	
		Percent	SE	Percent	SE	Percent	SE	Percent	SE
Hispanic	15,133	19.9	1.79	7.9	0.88	10.5	1.20	10.3	1.22
Non-Hispanic white	40,420	18.5	1.30	17.8	1.26	7.0	0.69	10.9	0.89
Non-Hispanic black	9,805	17.6	1.65	8.1	1.19	11.0	1.47	10.9	1.73

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. SE is standard error.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.

Data Brief 388. Current Contraceptive Status Among Women Aged 15–49: United States, 2017–2019

Data table for Figure 5. Percentage of all women aged 22–49 who were currently using female sterilization, oral contraceptive pills, the male condom, or long-acting reversible contraceptives, by educational attainment: United States, 2017–2019

Education	Number (thousands)	Female sterilization		Pill		Male condom		Long-acting reversible contraceptives	
		Percent	SE	Percent	SE	Percent	SE	Percent	SE
No high school diploma or GED	4,744	39.9	3.64	5.7	1.57	8.7	1.83	9.3	1.98
High school diploma or GED	14,530	31.5	2.30	6.8	1.28	9.3	1.17	7.9	0.84
Some college, but no bachelor's degree	17,631	22.2	1.56	10.8	1.24	8.7	1.05	12.5	1.08
Bachelor's degree or higher	22,371	12.1	1.36	18.1	1.57	8.6	1.03	13.1	1.39

NOTES: Women currently using more than one method are classified according to the most effective method they are using. Long-acting reversible contraceptives include contraceptive implants and intrauterine devices. SE is standard error.

SOURCE: National Center for Health Statistics, National Survey of Family Growth, 2017–2019.